

Exact testing for model/data fit for log-linear models

“Part Two”

“Algebraic & Geometric Methods in Statistics”

Sonja Petrović
Created for Math/Stat 561

Feb 27, 2026.

Agenda

- Chapter 9 from our textbook: Fisher's exact test
- Part of chapter 8, as we may need the cone of sufficient statistics.

Goals

- LAST LECTURE:
 - Understand hypotheses testing for model/data fit
- THIS LECTURE: we will work towards
 - What is a p -value for a goodness-of-fit test?
 - Asymptotic vs. exact tests
 - Fisher's test and example
- To ponder:
 - General goodness of fit test for log-linear models
 - Open problems and relation to projects!

Recap

Exact test (Fisher)

In an **exact** goodness-of-fit test, one uses the exact distribution of the statistic. . .

Recap

Exact test (Fisher)

In an **exact** goodness-of-fit test, one uses the exact distribution of the statistic... which is **what**?

	gender		
range	M	F	Nb
<=135K	8	1	4
> 135K	2	9	2

	gender		
range	M	F	Nb
<=135K	9	0	4
> 135K	1	10	2

	gender		
range	M	F	Nb
<=135K	9	1	3
> 135K	1	9	3

Conclusion? Evidence in the data? Significance?

Definition [p-value]

Refer to Chapter 5. Discuss in lecture / board.

- Read the beginning of Chapter 9. Section 9.1: Conditional inference.
 - We are *conditioning* on the row and column sums of the table.
 - These are sufficient statistics for the independence model.
 - This is a *general strategy*...

The general exact test for contingency tables [board lecture]

- Proposition 9.1.1. [stated without proof, but it's not difficult. . .]
- p.192 "A similar strategy is based on the likelihood ratio test, where we use the G statistic, instead of the X² statistic."
- Look back to the example from Lecture 10:

Interpret: what are all the possible tables? What is the probability of any given table?

	M	F	T/Nb	totals
$\leq 135K$?	?	?	13
$> 135K$?	?	?	13
totals	10	10	6	26

Here's a cheat sheet:

Before we proceed with the Fisher test, we first introduce some notations. We represent the cells by the letters a , b , c and d , call the totals across rows and columns *marginal totals*, and represent the grand total by n . So the table now looks like this:

	Men	Women	Row Total
Studying	a	b	$a + b$
Non-studying	c	d	$c + d$
Column Total	$a + c$	$b + d$	$a + b + c + d (=n)$

Fisher showed that conditional on the margins of the table, a is distributed as a [hypergeometric distribution](#) with $a+c$ draws from a population with $a+b$ successes and $c+d$ failures. The probability of obtaining such set of values is given by:

$$p = \frac{\binom{a+b}{a} \binom{c+d}{c}}{\binom{n}{a+c}} = \frac{\binom{a+b}{b} \binom{c+d}{d}}{\binom{n}{b+d}} = \frac{(a+b)! (c+d)! (a+c)! (b+d)!}{a! b! c! d! n!}$$

where $\binom{n}{k}$ is the [binomial coefficient](#) and the symbol ! indicates the [factorial operator](#). This can be seen as follows. If the marginal totals (i.e. $a+b$, $c+d$, $a+c$, and $b+d$) are known, only a single degree of freedom is left: the value e.g. of a suffices to deduce the other values. Now, $p = p(a)$ is the probability that a elements are positive in a random selection (without replacement) of $a+c$ elements from a larger set containing n elements in total out of which $a+b$ are positive, which is precisely the definition of the hypergeometric distribution.

Figure 1: From Wikipedia :)

Recap

Recap: definition of exact conditional tests

In an **exact goodness-of-fit test**, one uses the exact distribution of a GoF statistic. . . for example the X^2 statistic defined in lecture 10. The **reference distribution** is the *exact* distribution of that statistic on the set of tables with given fixed sufficient statistics (margins).

	gender		
range	M	F	Nb
<=135K	8	1	4
> 135K	2	9	2

	gender		
range	M	F	Nb
<=135K	9	0	4
> 135K	1	10	2

	gender		
range	M	F	Nb
<=135K	9	1	3
> 135K	1	9	3

Conclusion? Evidence in the data? Significance?

Definition [p-value]

Define the p -value for the GoF statistic.

- Read the beginning of Chapter 9. Section 9.1: Conditional inference.
 - We are *conditioning* on the row and column sums of the table.
 - These are sufficient statistics for the independence model.
 - This is a *general strategy*...

Interpret: what are all the possible tables? What is the probability of any given table?

	M	F	T/Nb	totals
$\leq 135K$?	?	?	13
$> 135K$?	?	?	13
totals	10	10	6	26

The general exact test for contingency tables [board lecture]

- Definition 9.1.3. - fiber
- p194: Problem 9.1.6. - understand the problem definition

Models with a design matrix

- X_1, \dots, X_k discrete random variables, $X_i \in \{1, \dots, d_i\}$
- $u =$ a **k-way contingency table** $u \in \mathbb{Z}_{\geq 0}^{d_1 \times \dots \times d_k}$ [Draw a table!] Flatten u to vector.

Log-linear model

Sufficient statistics = **marginals** of u : $P_\theta(U = u) = \exp\{\langle Au, \theta \rangle - \psi(\theta)\}$.

Example $X_1 \perp\!\!\!\perp X_2$

$$\left[\begin{array}{cccc|cccc|ccc}
 1 & 1 & \dots & 1 & 0 & 0 & \dots & 0 & \dots & 0 & 0 & \dots & 0 \\
 0 & 0 & \dots & 0 & 1 & 1 & \dots & 1 & \dots & 0 & 0 & \dots & 0 \\
 \vdots & \vdots \\
 0 & 0 & \dots & 0 & 0 & 0 & \dots & 0 & \dots & 1 & 1 & \dots & 1
 \end{array} \right]$$

$$\left[\begin{array}{cccc|cccc|ccc}
 1 & 0 & \dots & 0 & 1 & 0 & \dots & 0 & \dots & 1 & 0 & \dots & 0 \\
 0 & 1 & \dots & 0 & 0 & 1 & \dots & 0 & \dots & 0 & 1 & \dots & 0 \\
 \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \ddots & \vdots & \vdots & \vdots & \vdots & \ddots & \vdots \\
 0 & 0 & \dots & 1 & 0 & 0 & \dots & 1 & \dots & 0 & 0 & \dots & 1
 \end{array} \right]_{(d_1+d_2) \times d_1 d_2}$$

$$\begin{bmatrix} u_{11} \\ \vdots \\ u_{d_1 d_2} \end{bmatrix} = \begin{bmatrix} u_{1+} & \dots & u_{+d_2} \end{bmatrix}$$

Resources & License

- Quick summary [notes](#) about p -values that I wrote for Stat 514.
- Read about hypothesis tests for context of the model fitting tests in [these lecture notes](#).
- [This lesson](#) from Penn State online offers a one-page summary of Fisher's exact test for 2×2 tables, as it was developed by Sir Fisher!
- Believe it or not, there is a great 2×2 example on [Wikipedia](#), a page which actually contains a really good explanation for this one example.

This document is created for Math/Stat 561, Spring 2023.

All materials posted on this page are licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License](#).